

A stack of several books with blue covers and white pages, positioned on the left side of the slide. The books are slightly offset, showing the edges of the pages.

Chapter 28 Magnetic Force – Lorentz Force Law

Physics II – Part II
Wen-Bin Jian

Department of Electrophysics, National Chiao Tung University

Lorentz Force

History

1865 AD – James Clerk Maxwell (Scottish)
formulations of field equations

1881 AD – Sir Joseph John Thomson (English)
derive from Maxwell's equations

$$\vec{F} = \frac{1}{2} q \vec{v} \times \vec{B}$$

1881 AD – Oliver Heaviside (English)
derive the correct form of the force law


1892 AD – Hendrik Antoon Lorentz (Dutch)
modern form, named Lorentz force law due to
the consideration of relative transformation

$$\vec{F} = q\vec{E} + q\vec{v} \times \vec{B}$$

Is Lorentz law incompatible with special relativity?

<https://phys.org/news/2012-05-classical-electrodynamics-law-incompatible-special.html>

Masud Mansuripur, Phys. Rev. Lett. 108, 193901 (2012).



<http://en.citizendium.org/images/thumb/4/40/HALorentz.gif/250px-HALorentz.gif>

Hall Effect, Quantum Hall Effect, Fractional Quantum Hall Effect


History

1879 AD – Edwin Herbert Hall (American) an experimentalist for his doctoral degree at Johns Hopkins University in Baltimore, Maryland

1978 AD – Klaus von Klitzing (German) discover the **quantum Hall effect**

1982 AD – Robert B. Laughlin, Daniel C. Tsui, Horst L. Störmer (American) Laughlin, a theorist, explained the experimental results of **fractional quantum Hall effect** done by Tsui & Störmer.


<https://www.britannica.com/biography/Robert-B-Laughlin>



Physics Today 38(12), 17 (1985).

Magnetic Fields and Magnetic Forces

Lorentz Force Law



The Ampere's repulsive force between two opposite current flows

Alternative description for the repulsive force between two magnets

The mechanism could be separated to the interaction between the current and the magnetic force.

The Lorentz force law describes: charge in motion is exerted by a force from magnetic field

$$\vec{F}_{Lorentz} = q\vec{v} \times \vec{B}$$

Magnetic Fields

Lorentz Force Law

Source of Magnetic Field	Magnitude of Magnetic Field (T)
Transient Superconducting Magnet	100
Static Lab Superconducting Magnet	20
Conventional Electromagnet	2
Conventional Bar Magnet	0.01-1.5 (short range)
Medical MRI	1.5
Magnetic Field on The Earth	0.44×10^{-4} (0.5 Gauss)
Human Brain	10^{-15}

Static charges do not response to magnetic fields.

Magnetic force is perpendicular to both the magnetic field and the charge motion direction thus producing cycloid motion.

The Lorenz force law is a relative transformation of the static Coulomb law.

Moving Charges in Magnetic Fields

Cycloid Motion

Centripetal force is the driving force to make a circular motion.
Lorentz force is the centripetal force for moving charges.



Cyclotron frequency is independent of the velocity of the charged particle.

$$F = qvB = ma_r \quad \vec{v} \perp \vec{B}$$

$$q(r\omega)B = m(r\omega^2) \rightarrow \omega = \frac{qB}{m}$$

only depends on the charge, the mass, and the magnetic field

If the motional particle possess a velocity component parallel to the magnetic field, it will undergoes a constant velocity motion in that direction.




Instrumentation

Application of Lorenz Force

The velocity selector. Use magnetic and electric fields to select the charged particles with specified velocities.

$$qE = qvB \rightarrow v = \frac{E}{B}$$




Application of
Lorenz Force

Mass spectrometer

$$qvB = \frac{mv^2}{r} \rightarrow v = \frac{qBr}{m}$$

$$K = qV = \frac{1}{2}mv^2 = \frac{1}{2}m \left(\frac{qBr}{m} \right)^2$$

$$V = \frac{1}{2} \frac{qB^2 r^2}{m} \rightarrow m = \frac{qB^2 r^2}{2V}$$




Cyclotron accelerator

$$\omega = \frac{qB}{m}$$

Large radius gives high energy for the charged particles.

$$K = \frac{1}{2}m(\omega R)^2 = \frac{q^2 B^2}{2m} R^2$$



Another Form of Lorentz Force Law

Lorentz Force on Current-Carrying Wires

In the segment of a conductor, there are N charged particles

$$N = nLA$$

$$\vec{F}_{net} = Nq\vec{v} \times \vec{B} = lAnq\vec{v} \times \vec{B}$$

$$\vec{F}_{net} = lA\vec{J} \times \vec{B} = I\vec{l} \times \vec{B}$$

$$d\vec{F} = I(d\vec{l}) \times \vec{B} \longrightarrow \vec{F}_{net} = I \int d\vec{l} \times \vec{B}$$



Torque on the current loop:

$$F = IabB$$

$$\tau = \frac{b}{2}(IaB) \sin(\theta) \times 2$$

$$\vec{\tau} = (Iab)B \sin(\theta) = (Iab\hat{n}) \times \vec{B}$$

Define magnetic moment $\vec{m} = Iab\hat{n}$



Lorentz Force on a Current Loop, Magnetic Moment, Torque, Potential Energy

$$\vec{\tau} = \vec{m} \times \vec{B}$$

Potential energy: exerted external torque


$$\tau_{ext} = -\tau = -mB \sin(\theta)$$

$$dW = (-mB \sin(\theta))d\theta$$

$$dU = mB \sin(\theta) d\theta$$


$$U = \int_{\pi/2}^{\theta} mB \sin(\theta) d\theta = -mB \cos(\theta)$$

$$U = -\vec{m} \cdot \vec{B}$$



Torque on a Current Loop

Determine The Polarity of Charged Particles and The Carrier Concentration



The Hall effect induced electric field:

$$F = qv_d B = qE_H \rightarrow E_H = v_d B$$

The Hall voltage is $V_H = v_d B w$

$$I = (wt)nqv_d \rightarrow v_d = I/wtnq \quad V_H = \frac{IBw}{wt nq} = \frac{IB}{nqt}$$

$$V_H = IR \rightarrow R = \frac{B}{nqt} = \frac{1}{nq} \frac{B}{t} = R_H \frac{B}{t} \quad R_H = \frac{1}{nq}$$

$$n = \frac{IB}{qtV_H} \quad \text{Quantum Hall: } R = \frac{V_H}{I} \rightarrow G = \frac{I}{V_H} = \frac{nqt}{B} = m\left(\frac{e^2}{h}\right)$$

Hall Effect

Lorentz Force on a Segment of Current-Carrying Conductor

A proton of mass $m = 1.67 \times 10^{-27}$ kg and charge $q = e = 1.602 \times 10^{-19}$ C moves in a circle of radius $r = 21$ cm perpendicular to a magnetic field of $B = 4000$ G. Find (a) the period of the motion and (b) the speed of the proton.

$$qvB = m \frac{v^2}{r} \rightarrow v = \frac{qBr}{m} = \frac{(1.602 \times 10^{-19})(0.4)(0.21)}{(1.67 \times 10^{-27})}$$

$$v = 8.05 \times 10^6 \text{ m/s}$$

$$T = \frac{2\pi r}{v} = \frac{2(3.14159)(0.21)}{8.05 \times 10^6} = 1.64 \times 10^{-7} \text{ s}$$

Examples

The Mass Spectrometer

Examples

A ^{58}Ni ion of charge $+e$ and mass 9.62×10^{-26} kg is accelerated through a potential drop of 3 kV and deflected in a magnetic field of 0.12 T. (a) Find the radius of curvature of the orbit of the ion. (b) Find the difference in the radii of curvature of ^{58}Ni ions and ^{60}Ni ions.

$$(a) \quad qV = \frac{1}{2}mv^2 \quad \& \quad \omega = \frac{qB}{m} \rightarrow qV = \frac{1}{2}mr^2 \frac{q^2B^2}{m^2}$$
$$r_{^{58}\text{Ni}} = \sqrt{2mV/qB^2} = \sqrt{\frac{2 \times (9.62 \times 10^{-26}) \times (3000)}{(1.602 \times 10^{-19}) \times (0.12)^2}} = 0.501 \text{ m}$$

$$(b) \quad \frac{r_{^{60}\text{Ni}}}{r_{^{58}\text{Ni}}} = \sqrt{\frac{60}{58}} \rightarrow r_{^{60}\text{Ni}} = \sqrt{\frac{60}{58}} \times 0.501 = 0.510 \text{ m}$$

$$\Delta r = 0.009 \text{ m}$$

The Cyclotron

Examples

A cyclotron for accelerating protons has a magnetic field of 1.5 T and a maximum radius of 0.5 m. (a) What's the cyclotron frequency? (b) What's the kinetic energy of the protons when they emerge?

$$(a) \quad \omega = \frac{qB}{m} = \frac{(1.602 \times 10^{-19}) \times (1.5)}{1.67 \times 10^{-27}} = 1.44 \times 10^8 \text{ rad/s}$$

$$f = \frac{\omega}{2\pi} = 2.29 \times 10^7 \text{ Hz}$$

$$(b) \quad K = \frac{1}{2}mv^2 = \frac{1}{2}m\omega^2r^2 = 4.33 \times 10^{-12} \text{ J} = 27.0 \text{ MeV}$$

Lorentz Force on a Segment of Current-Carrying Conductor


A wire segment of 1 mm in length carries a current of 1.0 A in the +x direction. If a magnetic field of 1.0 T is applied perpendicular to the z axis and making an angle of $\pi/4$ with the x-axis. What is the magnitude of the Lorentz force?

$$F = ILB \sin(\theta)$$

$$F = (1)(0.001)(1) \sin(\pi/4)$$

$$F = 7.1 \times 10^{-4} \text{ N}$$

$$\vec{F} = 7.1 \times 10^{-4} \hat{k} \text{ N}$$



Examples

Lorentz Force on a Curved Wire of Current-Carrying Conductor

A wire bent into a semicircular loop of radius R lies in the xy plane and carries a current I . It is placed in a field $\vec{B} = B_0 \hat{k}$. Please find the Lorentz force exerted on the wire.


$$dF = IdlB_0$$

$$dl = Rd\theta$$

$$F_x = 0, F_y = \int dF_y$$

$$dF_y = I(Rd\theta)B_0 \sin(\theta)$$

$$F_y = \int_0^\pi IRB_0 \sin(\theta) d\theta = 2IRB_0$$



Examples


Magnetic Torque on a Current Loop

A circular loop of wire with a radius R and mass m carries a current I and lies in the horizontal plane. A horizontal magnetic field B is applied in the space. How large can the current be before one edge of the loop is lifted off the plane?

$$m_B = IA = I\pi R^2$$

$$\tau_B < \tau_g \rightarrow I\pi R^2 B < mgR$$

$$I < \frac{mg}{\pi RB}$$



Examples

Magnetic Moment & Torque on a Current Loop

Examples

A square 12-turn coil with edge-length 40 cm carries a current of 3 A. It lies in the xy plane in a uniform magnetic field $\vec{B} = 0.3\hat{i} + 0.4\hat{k}$ (T). Find (a) the magnetic moment of the coil, (b) torque, and (c) the potential energy.

(a)

$$\vec{m} = 12 \times 3 \times (0.4)^2 \hat{k} = 5.76 \hat{k} \text{ (Am}^2\text{)}$$


(b)

$$\vec{\tau} = \vec{m} \times \vec{B} = (5.76 \hat{k}) \times (0.3\hat{i} + 0.4\hat{k})$$

$$\vec{\tau} = 1.73 \hat{j}$$

(c)

$$U = -\vec{m} \cdot \vec{B} = -(5.76 \hat{k}) \cdot (0.3\hat{i} + 0.4\hat{k}) = 2.30 \text{ J}$$



Calculation of Magnetic Moment


A thin nonconducting disk of mass m and radius R has a uniform surface charge per unit area σ and rotates with angular velocity ω about the axis. Find the magnetic moment.

$$\mu = IA$$

$$d\mu = AdI \rightarrow A = \pi r^2, dI = \frac{dQ}{T} = \frac{\sigma 2\pi r dr}{2\pi/\omega}$$

$$d\mu = (\pi r^2)(\sigma \omega r dr)$$

$$\mu = \int_0^R \sigma \omega \pi r^3 dr = \frac{\sigma \omega \pi}{4} R^4$$



Examples